



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

2690 [April, 1918]. Proposed by E. V. HUNTINGTON, Harvard University.

Find the maximum value of

$$y = \frac{\sin x \cos (x + \varphi)}{\cos (x + \theta) \sin (x + \beta + \theta)}. \quad (1)$$

SOLUTION BY J. B. REYNOLDS, Lehigh University.

Taking the logarithm of both sides, differentiating, and setting the result equal to zero we get

$$\cot x - \tan (x + \varphi) + \tan (x + \theta) - \cot (x + \beta + \theta) = 0 \quad (2)$$

and from the two forms of this

$$\cot x - \cot (x + \beta + \theta) = \tan (x + \varphi) - \tan (x + \theta)$$

and

$$\cot x + \tan (x + \theta) = \cot (x + \beta + \theta) + \tan (x + \varphi)$$

we get

$$\frac{\sin (\beta + \theta)}{\sin (\varphi - \theta)} = \frac{\sin x \sin (x + \beta + \theta)}{\cos (x + \varphi) \cos (x + \theta)} \quad (3)$$

and

$$\frac{\cos (x + \varphi)}{\cos (x + \theta)} = \frac{\cos (\beta + \theta - \varphi)}{\cos \theta} \cdot \frac{\sin x}{\sin (x + \beta + \theta)}. \quad (4)$$

By (3) and (4), we have

$$\sqrt{\frac{\cos (\beta + \theta - \varphi) \sin (\varphi - \theta)}{\cos \theta \sin (\beta + \theta)}} = \frac{\cos (x + \varphi)}{\sin x} = \cot x \cos \varphi - \sin \varphi.$$

Hence,

$$\cot x = \tan \varphi + \sec \varphi \sqrt{\frac{\cos (\beta + \theta - \varphi) \sin (\varphi - \theta)}{\cos \theta \sin (\beta + \theta)}}. \quad (5)$$

(4) and (1) give, where y_1 is the maximum value of y ,

$$y_1 = \frac{\cos (\beta + \theta - \varphi)}{\cos \theta} \cdot \frac{\sin^2 x}{\sin^2 (x + \beta + \theta)};$$

whence

$$\frac{1}{\sqrt{y_1}} = \sqrt{\frac{\cos \theta}{\cos (\beta + \theta - \varphi)}} \{ \cos (\beta + \theta) + \sin (\beta + \theta) \cot x \}$$

and eliminating $\cot x$ by (5), we have

$$y_1 = \cos^2 \varphi / \{ \sqrt{\cos \theta \cos (\beta + \theta - \varphi)} + \sqrt{\sin (\varphi - \theta) \sin (\beta + \theta)} \}^2.$$

As in 2689 we may prove that y_1 is a maximum as found for the case $\beta = \varphi$ from which we may infer from the manner that β enters into the expression that it is a maximum for all values of β .

2692 [April, 1918]. Proposed by J. L. RILEY, Stephenville, Texas.

A cube is cut at random by a plane, what is the chance that the section is a hexagon?

SOLUTION BY C. F. GUMMER, Queen's University, Kingston, Ont.

Let the faces of the cube be $x = \pm 1$, $y = \pm 1$, and $z = \pm 1$. Let the plane be

$$lx + my + nz = p,$$

where $n = \cos \theta > 0$, $l = \sin \theta \cos \phi$, $m = \sin \theta \sin \phi$. The normal from the origin is equally likely to have any direction, and, when the direction is fixed, equally likely to have any length. Since $\sin \theta d\theta d\phi$ is an element of the solid angle traced by the normal, and dp an element of its length, the relative probability of the plane lying within given limits is $\iiint \sin \theta dp d\theta d\phi$. On account of the symmetry of the cube, it may be assumed that the normal through O to the plane meets the surface of the cube within the triangle $(0, 0, 1)$, $(1, 0, 1)$, and $(1, 1, 1)$; so that $n > l > m > 0$, that is, $0 < \phi < \pi/4$, $0 < \theta < \tan^{-1} \sec \phi$.